

Total hip arthroplasty improves pain and function but not physical activity

Jeldi, Artaban Johnson; Deakin, Angela H.; Allen, David J.; Granat, Malcolm H.; Grant, Margaret; Stansfield, Ben W.

Published in:
Journal of Arthroplasty

DOI:
[10.1016/j.arth.2017.02.002](https://doi.org/10.1016/j.arth.2017.02.002)

Publication date:
2017

Document Version
Author accepted manuscript

[Link to publication in ResearchOnline](#)

Citation for published version (Harvard):
Jeldi, AJ, Deakin, AH, Allen, DJ, Granat, MH, Grant, M & Stansfield, BW 2017, 'Total hip arthroplasty improves pain and function but not physical activity', *Journal of Arthroplasty*, vol. 32, no. 7, pp. 2191-2198.
<https://doi.org/10.1016/j.arth.2017.02.002>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

Title: Total hip arthroplasty improves pain and function but not physical activity

Artaban Johnson Jeldi^{a,b}, PhD (artaban.jeldi@gjnh.scot.nhs.uk)

Angela.H. Deakin^b PhD (Angela.Deakin@gjnh.scot.nhs.uk)

David Allen^b, FRCSEd (Trauma & Orthopaedics) (David.Allen@gjnh.scot.nhs.uk)

Malcolm. H. Granat^c PhD (m.h.granat@salford.ac.uk)

Margaret Grant^a, PhD (M.grant@gcu.ac.uk)

Ben Stansfield^a PhD (ben.stansfield@gcu.ac.uk)

Authors: Address

- a) Institute for Applied Health Research, Glasgow Caledonian University, Glasgow,
UK G4 0BA.
- b) Department of Orthopaedics, Golden Jubilee National Hospital, Clydebank, Glasgow,
UK. G81 4DY.
- c) School of Health Sciences, University of Salford, Salford, UK. M6 6PU.

Corresponding Author

Dr Artaban Jeldi,

Orthopaedics Research, R&D Department, Golden Jubilee National Hospital, Clydebank,
Glasgow, UK

E-mail: artaban.jeldi@gjnh.scot.nhs.uk, ben.stansfield@gcu.ac.uk

Abstract

Background:

People with hip osteoarthritis are likely to limited physical activity (PA) engagement due to pain and lack of function. Total hip arthroplasty (THA) reduces pain and improves function, potentially allowing increased PA. PA of THA patients was quantified to 12m post-operation. The hypothesis was that post-operatively levels of PA would increase.

Methods:

PA of 30 THA patients (67 ± 7 years) was objectively measured pre-operatively and three and 12 months post-operation. Harris Hip Score (HHS), Oxford Hip Score (OHS) and six minute walk test (6MWT) were recorded. Mixed linear modelling was used to examine relationships of outcomes with time, BMI, age, gender and baseline HHS.

Results:

Time was not a significant factor in predicting sit-to-stand transitions, upright time, steps, cadence of walking bouts >60s, or longest upright bouts. However, HHS, OHS and 6MWT all improved with time. Notably BMI was a significant predictor of upright time, steps, largest number of steps in an upright bout, HHS and 6MWT. Baseline HHS helped predict longest upright bout, cadence of walking bouts >60s and OHS. The significant effect of participant as a random intercept in the model for PA outcomes suggested habituation from pre- to post-surgery.

Conclusions:

PA did not change from pre- to 12m post-surgery despite improvement in HHS, OHS and 6MWT. Baseline BMI was a more important predictor of upright activity and stepping than time. Pre- and post-operative PA promotion could be used to modify apparently habitual low levels of PA to enable full health benefits of THA to be gained.

Title: Total hip arthroplasty improves pain and function but not physical activity

Abstract

Background:

People with hip osteoarthritis are likely to limited physical activity (PA) engagement due to pain and lack of function. Total hip arthroplasty (THA) reduces pain and improves function, potentially allowing increased PA. PA of THA patients was quantified to 12m post-operation. The hypothesis was that post-operatively levels of PA would increase.

Methods:

PA of 30 THA patients (67 ± 7 years) was objectively measured pre-operatively and three and 12 months post-operation. Harris Hip Score (HHS), Oxford Hip Score (OHS) and six minute walk test (6MWT) were recorded. Mixed linear modelling was used to examine relationships of outcomes with time, BMI, age, gender and baseline HHS.

Results:

Time was not a significant factor in predicting sit-to-stand transitions, upright time, steps, cadence of walking bouts >60 s, or longest upright bouts. However, HHS, OHS and 6MWT all improved with time. Notably BMI was a significant predictor of upright time, steps, largest number of steps in an upright bout, HHS and 6MWT. Baseline HHS helped predict longest upright bout, cadence of walking bouts >60 s and OHS. The significant effect of participant as a random intercept in the model for PA outcomes suggested habituation from pre- to post-surgery.

Conclusions:

PA did not change from pre- to 12m post-surgery despite improvement in HHS, OHS and 6MWT. Baseline BMI was a more important predictor of upright activity and stepping than

time. Pre- and post-operative PA promotion could be used to modify apparently habitual low levels of PA to enable full health benefits of THA to be gained.

Keywords

Physical activity, Total Hip Arthroplasty

List of Abbreviations

THA Total hip arthroplasty

PA Physical activity

HHS Harris Hip Score

OHS Oxford Hip Score

6MWT Six minute walk test

BMI Body mass index

41 Introduction

42 Total hip arthroplasty (THA) leads to reduction in pain[1–3] and improvement in functional
43 capacity[4–6] and quality of life[7,8]. Additionally THA is associated with enhanced walking
44 endurance (longer distances covered in the 6 minute walk test[9,10]) and improved balance
45 (faster performance of the timed-up-and-go-test[10,11]). However, there is emerging
46 evidence that there are only small changes in free-living physical activity (PA) following
47 surgery[12–15]. With reduction in pain, enhancement of endurance and speed of walking
48 following THA, it might be expected that individuals would be able to participate in health
49 enhancing PA. However, it is possible that pre-operative habitual patterns of PA are not
50 altered. If pre-operative patterns of PA are ingrained (at low levels), then pre-operative
51 measurements could be extremely useful in targeting person-centred interventions to
52 disrupt these patterns, potentially enhancing long-term health prospects. While PA may
53 have become habituated, it is possible that changes following surgery may be dependent on
54 a range of factors. A person's gender may have an impact on surgical outcomes, as might
55 their weight and age at the point of surgery. Also their pre-operative clinical condition may
56 be important in determining outcomes.

57 The aim of this study was to enhance understanding of PA following THA by using objective
58 measurement to characterise PA from pre-operation through recovery to twelve months
59 post-operation. The relationship between PA change with time following surgery and
60 gender, age, BMI and baseline clinical score (Harris Hip Score) was investigated. Primary
61 physical activity outcomes included the number of sit-to-stand transitions per day, the time
62 upright per day and the number of steps per day. Also secondary outcomes were quantified
63 to further characterise PA (characteristics of the longest bouts of activity), walking

endurance (six minute walk test) and clinical outcomes (Harris Hip Score, Oxford Hip Score).
The hypothesis was that all outcomes would improve following THA.

Patients and Methods

This was an observational cohort study. Ethical approval was obtained from the West of Scotland Research Ethics Committee 1 (12/WS/0098). The study population was all patients being seen for a primary total hip arthroplasty operation at an NHS elective arthroplasty centre. To ensure the external validity of the study results the inclusion / exclusion criteria were kept as wide as possible. Inclusion criteria for the study were patients who were able to give informed consent, were between 50-85 years old and could return for follow-up. Patients were excluded if they were undergoing a revision hip arthroplasty, had had a total hip or knee arthroplasty in the last 12 months, had extreme locomotor limitations due to cardio-pulmonary, central or peripheral nervous system deficits or spinal conditions or were diagnosed with a terminal disease (malignancy). From July to August 2012 and January to May 2013 (break in recruitment due to illness of lead investigator) all THA patients under the care of one consultant orthopaedic surgeon (n=64) were reviewed for eligibility for the study. All eligible patients (n=57) were approached for inclusion in the study. Published data to carry out an appropriate a-priori power calculation were not available. Therefore a target sample of 30 participants was set to provide a power of 0.8 to detect a difference of 1SD in outcomes with a significant level of 0.05. All the study participants gave informed consent and the complete assessment was carried out by the lead author, a registered Physiotherapist.

Participants were operated on by a single consultant surgeon (DA) or a trainee surgeon under his direct supervision. Using a posterior approach, all participants received either a Contemporary® cemented cup or Trident® uncemented cup with an X3 polyethylene liner and an Exeter® femoral component (Stryker Orthopaedics, Michigan, USA). The peri-operative care for all participants (from pre-assessment through to discharge) followed the institution's enhanced recovery programme[16]. The aim of this programme was to accelerate patients' rehabilitation and reduce in-hospital length of stay (to 3-4 days) by implementing a multimodal anaesthetic regime combined with pre-operative education and early mobilisation.

Data were collected at three time points: Pre-operatively within the two weeks before operation; three months after operation; twelve months after operation.

PA data were collected objectively for up to seven days using the activPAL3™ monitor (50x35x7mm, 30g) (PAL Technologies Ltd. Glasgow, UK; software version 7.1.18)[17,18].

Data from this instrument classified activities into sedentary (sitting/lying), standing and stride events. Thus a record of posture (upright or not) and stepping activity was generated.

The monitor was attached to the anterior aspect of the thigh of the non-operated leg (24 hour/day wear) using a waterproof surgical dressing (Duoderm extra thin hydrocolloid dressing (Convatec) or Opsite flexifix (Smith & Nephew)). When compared to video based observation the activPAL3™ has only a 0.27% difference in upright time detection and -3.34% difference for step count in adults [19] during standardised activities. Whilst the upright time detection remains good in activities of daily living (-0.19% agreement) there is considerable undercounting of small stepping activity within these activities (-86%). This undercounting of small/slow steps associated with some ADLs is emphasised by the

109 monitor's progressively poorer performance below 0.5m/s [20]. The monitor therefore
110 records purposeful stepping activity, but is poor at recording small/slow incidental stepping.

111 Compliance with monitor wear was assessed by self-report and post-hoc data examination
112 by the research team. Only self-reported full 24 hours periods of wear were considered for
113 inclusion. These records were manually inspected for apparent abnormalities. Any days with
114 apparent abnormalities were excluded.

115

116 *Primary outcome measures:*

117 The primary outcome for the study was PA. To quantify the level of PA the following three
118 outcome measures were averaged over all 24 hour periods recorded: Sit-to-stand
119 transitions/day; time spent upright/day (hours/day); steps/day.

120 *Secondary outcome measures:*

121 The following characteristics of the longer bouts of activity for each recording period were
122 calculated: Duration of longest continuous upright bout over the recording period; largest
123 number of steps in an upright bout across the recording period (not necessarily the same
124 bout as the longest continuous upright bout); mean cadence (steps/min) of all walking bouts
125 of longer than 60s.

126 Participants were also assessed at all time points using the Harris Hip Score[21] (completed
127 by the lead author by measurement and interview), the Oxford Hip Score[22,23] (self-
128 completion) and a six minute walk test [24]. The six minute walk test was conducted in a
129 30m long corridor, which had regular rest stations at 10m intervals. The participants were

advised to walk up and down the corridor at a self-selected speed to achieve the maximum distance within 6 minutes. The participants were informed that they could take a break when and for how long needed at any point during the 6 minutes.

The age, gender, height, weight, BMI and comorbidities (from medical notes) of participants were recorded. Major complications (Death, Pulmonary Embolism/Deep vein Thrombosis, dislocation, infection, and revision) were noted.

Data analysis and statistics

Outcomes were characterised as mean (SD). To examine the change pre- to post-operation mixed linear models were used to model the relationship between each PA or clinical outcome parameter and time. As three time points were recorded time could only be modelled as quadratic. Models were adjusted for gender, baseline BMI, age and baseline Harris Hip Score (apart from Harris Hip Score). The shape of each outcome curve over time was modelled with each outcome measurement at Level 1 and each patient at Level 2. Fixed and random effects were included at the patient level (Level 2) and measurement level (Level 1). The fixed part of each model describes the average growth curve for the sample; the random part splits variation between subjects at the higher level and variation between time in the study of the same person at the lower level. The models allowed a unique growth curve to be generated for each subject based on his or her deviation from the average curve. An unstructured covariance structure was used.

Models were created adding covariates sequentially and comparing models using the $-2 \times \log \text{Likelihood}$ (-2LnL). Covariates were only retained if there was a significant improvement in -2LnL (reduction of 3.84). The maximum likelihood method was used to

152 estimate the coefficients so that unbiased estimates of $-2\ln L$ were calculated. Once the
153 model had been selected the restricted maximum likelihood method was used to give
154 unbiased estimates of the coefficients. From this model predictions were made. Residuals
155 for both level 2 and level 1 were estimated and investigated, as was the variance of the
156 model over time. Model parameters and 95%CI are presented along with predictions.

157 All statistical analysis was conducted in SPSS 23 (SPSS Inc, Chicago, IL) with the level of
158 statistical significance taken to be $p < 0.05$.

159

Results

Thirty participants (21F/9M) were recruited to the study. The STROBE flow diagram giving recruitment pathway and reasons for non-participation is given in Figure 1. All surgeries were successfully carried out. For the duration of the study (one year follow-up) there were no major complications. Of the 30 participants, 3 did not have 12 month data (Figure 1). All 30 participants' data were included in the mixed linear modelling. For the participants mean age was 67 years (range 50-82y), height 165cm (range 150-182cm), weight 82.9kg (range 57.8-132.6kg), BMI 31 kg/m² (range 19-43kg/m²) and pre-operative Harris Hip Score 50 (range 27-66) and Oxford Hip Score 15 (range 4-30). Indication for surgery in all the participants was osteoarthritis. Along with this diagnosis 13 participants had hypertension, 7 had cardiac abnormalities, 4 had Diabetes Mellitus, 7 had asthma/COPD and 2 participants had previous THA on the contralateral side. A median of six days of PA data were recorded at each time point. Mean and standard deviation of outcomes are presented (Table 1) with significant predictors within the mixed linear models (Table 2) and graphical evidence of trends with these significant predictors (Figure 2).

When examined using mixed linear models gender was not a significant predictor of any outcomes. Age was only a significant predictor of HHS ($b=0.38$, $p=0.047$) with a higher age being associated with a higher HHS (Figure 2g).

Time was not a significant predictor of the number of sit-to-stand transitions per day, upright time per day, steps per day, longest upright bout, the largest number of steps in an upright bout or cadence based on the outcomes of the mixed modelling (Table 1 mean data, Table 2 model outcomes). However, BMI was a significant predictor of upright time per day ($b=-0.153$, $p=-0.003$) (Figure 2a), steps per day ($b=-263$, $p=0.001$) (Figure 2b) and largest

number of steps in an upright bout ($b=-144$, $p=0.001$) (Figure 2d). In each of these cases higher BMI predicted a lower level of PA. Additionally Baseline HHS predicted variation in the longest upright bout ($b=0.0325$, $p=0.053$) (Figure 2c) and the cadence of walking in bouts longer than 60s ($b=0.696$, $p=0.004$) (Figure 2e). For all PA outcomes except the longest upright bout random intercepts explained a significant proportion of the outcome, indicating that participants tended to maintain the same level of PA in relation to the other participants across the study period.

Time and Time² were significant factors in the model for HHS, OHS and 6MWT outcomes (all $p<0.001$). Also BMI helped to predict outcomes for HHS ($b=-0.41$, $p=0.070$) (Figure 2f) and 6MWT ($b=-6.09$, $p=0.03$) (Figure 2i) with higher BMI indicating lower scores. HHS baseline value helped to describe OHS ($b=0.19$, $p=0.053$) (Figure 2h) change over the study period. There was significant variation across the study period between participants in HHS, OHS and 6MWT as indicated by significant (-2LnL) contributions to the model of random intercepts.

Discussion

This study demonstrated that at one year post-operation primary THA patients had made little change to their free-living PA from pre-operation levels, with time not being a predictor in the mixed linear model. However, BMI was a significant predictor in the model for upright time and stepping activity. The significance of the random parameter of participant within the model coupled with the lack of a significant effect of time, suggested that participants were tending to maintain the same relative volumes of PA across the study

period. This appears to indicate that pre-operative PA may have become habitual (possibly related to BMI) and despite improvements in function of the joint, as seen in the improvement of clinical outcome measures, PA levels did not significantly increase post-operation.

There were a number of limitations to this study. The small size of the study and the participants being under the care of one consultant within one hospital could limit the generalisability of the results. In terms of study power, based on the standard deviation of the difference in steps per day from pre- to 12 months post-operation recorded in this study (2492), the sample used (27 full records) would have been sufficient to detect a difference of 1350 steps/day with a power of 0.8 and a confidence of 0.05. In support of the generalizability of the results, the participants' age was similar to that reported in arthroplasty registers [25,26] and similarly osteoarthritis was the main reason for operation. However, BMI (mean 31kg/m²) was higher than reported elsewhere[25,27]. It is widely accepted that by 12 months post-operatively patients have gained the maximum benefit from their THA. However, it is possible that increased function is obtained at longer follow-up or that function at one year had already begun to deteriorate due to other co-morbidities. The original activPALTM monitor has proven validity in adults [18] and older adults[17,28]. However, the step counting facility of the activPAL3TM has limitations at slow stepping speeds as it under-counts slow, short step-length steps [20,29]. Therefore, the monitor may not have reliably detected stepping which was not 'purposeful'. A further issue is that this was an observational study of a surgical intervention. Theoretically, it would have been possible to perform an RCT with a no surgery arm to examine natural progression

within this population. A more subtle limitation is that this study did not measure the desire of participants to increase their level of PA post-surgery. Therefore, even with improved function participants may not have increased PA as they lacked motivation.

The results of this study confirm the previous reports of only small changes in PA following THA [14,15,30,31]. However, there were differences in outcomes, e.g. the current study (Table 1) found lower sit-to-stand transitions per day both pre-operatively and at 3 months post-operatively and whilst the upright time/day was similar pre-operatively, there were varying levels of agreement post-operatively compared to other studies. There were differences in participant demographics, with the current study population being older and more overweight than those of previous reports, which might explain these differences.

The mixed model outcomes indicated that gender did not significantly predict results for any outcome parameters within this cohort and that age was only significant within the prediction model for HHS. These results are perhaps surprising in that age might have been considered important in predicting PA as the study participants covered 50-82 years of age. However, BMI or Baseline HHS appeared to be more important. BMI was a predictor of both PA and clinical score outcomes, highlighting the importance of pre-operative BMI in predicting outcomes following surgery.

The inclusion of the Time² term within the mixed model improved predictions of several outcomes (Harris Hip Score, Oxford Hip Score and six minute walk test), suggesting that there was a non-linear relationship with time. This is highlighted in mean scores for these outcomes where large improvements occurred to the 3 month post-operative time point, but only small changes from 3 to 12 months post-operative (Figures 2f-i).

In the current study additional PA measures were added to those previously reported. The longest upright bout and largest number of steps in an upright bout provide quantification of the longest times participants performed 'functional' tasks requiring the upright posture. Time was not a significant factor in predicting these outcomes (Table 2), indicating that participants were not extending their loaded use of their new hip joints. Similarly cadence of stepping for bouts longer than 60s gives an insight into the intensity of stepping over extended periods. Whilst cadence increased across the study period (Table 1), this was not significantly predicted by time (Table 2) and still remained below that of age matched peers (93 (\pm 12) against 107 steps/min) [32–34]. The longest upright bout model did not have a significant random effect of participant suggesting that there was not a consistent ranking of participants with time (Table 2). It is possible that this outcome is highly influenced by particular social events or functional activities, pointing to a need to gather contextual information to gain a full understanding of the reasons for these patterns. However, the largest number of steps in an upright bout did have significant random effects of participant, suggesting similar volumes of stepping within one bout across the study period by participants.

The lack of time as a significant predictor within the model coupled with the significant random effect of participant suggests that pre-operative PA (except longest upright bout) may be habituated. Therefore, if pre-operative PA was measured, interventions could be used to target those likely to have low PA post-operatively to attempt to modify long-term behaviour. Enhancing PA has demonstrated secondary benefits of improving health (e.g. lower risk of cancer, ischemic events, diabetes [35] and enhanced quality of life[36,37]). A behaviour change intervention delivered through educational material, therapy sessions

etc., could be used to attempt to maximise the potential gains from THA in terms of overall health improvement. The significance of BMI within several PA outcome models reinforces the need to consider this as an important factor in health promotion alongside PA promotion.

As expected [9,30], both Harris Hip Score and Oxford Hip Score highlighted improvement from pre- to post-operation, as did 6MWT (values similar to previous studies [12,31]). However, there was not an accompanying increase in PA levels, indicating that these measures cannot be used as surrogates for PA, i.e. that it is necessary to measure free-living PA directly to gain insight into any changes following surgery.

Conclusions

In this study primary total hip arthroplasty patients did not make significant changes in the volume of PA performed at one year post-operation and it appeared that participants tended to maintain the same relative level of PA in relation to their peers. However, standard clinical outcome measures improved, showing an increase in function. This may indicate that habitual free-living PA patterns are established pre-operatively, perhaps related to BMI, and these are not altered by the better function and pain reduction given by a THA. These results may indicate that intervention to modify habitual low levels of PA, associated with declining long term health, could be necessary in a proportion of primary THA patients to allow them to fully exploit the additional function that their new joint gives them.

294 **References:**

- 295 [1] Montin L, Leino-Kilpi H, Suominen T, Lepistö J. A systematic review of empirical
296 studies between 1966 and 2005 of patient outcomes of total hip arthroplasty and
297 related factors. *J Clin Nurs* 2008;17:40–5. doi:10.1111/j.1365-2702.2007.01944.x.
- 298 [2] Banerjee M, Bouillon B, Banerjee C, Băthis H, Lefering R, Nardini M, et al. Sports
299 activity after total hip resurfacing. *Am J Sports Med* 2010;38:1229–36.
300 doi:10.1177/0363546509357609.
- 301 [3] Learmonth ID, Young C, Rorabeck C. The operation of the century: total hip
302 replacement. *Lancet* 2007;370:1508–19. doi:10.1016/S0140-6736(07)60457-7.
- 303 [4] Vissers MM, Bussmann JBJ, de Groot IB, Verhaar J a N, Reijman M. Walking and chair
304 rising performed in the daily life situation before and after total hip arthroplasty.
305 *Osteoarthritis Cartilage* 2011;19:1102–7. doi:10.1016/j.joca.2011.06.004.
- 306 [5] Bhav A, Mont M, Tennis S, Nickey M, Starr R, Etienne G. Functional problems and
307 treatment solutions after total hip and knee joint arthroplasty. *J Bone Joint Surg Am*
308 2005;87 Suppl 2:9–21. doi:10.2106/JBJS.E.00628.
- 309 [6] Dawson J, Fitzpatrick R, Murray D, Carr A. Comparison of measures to assess
310 outcomes in Total Hip Replacement surgery. *Qual Heal Care* 1996;5:81–8.
- 311 [7] Sliwinski M, Sisto S. Gait, quality of life, and their association following total hip
312 arthroplasty. *J Geriatr Phys Ther* 2006;29:10–7.
- 313 [8] Andreas L, Robert B. The Effect of of Elective Total Hip Replacement on Health-
314 Related Quality of Life. *J Bone Jt Surg* 1993;75:1619–25.
- 315 [9] Heiberg KE, Ekland A, Bruun-Olsen V, Mengshoel AM. Recovery and prediction of
316 physical functioning outcomes during the first year after total hip arthroplasty. *Arch*
317 *Phys Med Rehabil* 2013;94:1352–9. doi:10.1016/j.apmr.2013.01.017.
- 318 [10] Kennedy DM, Stratford PW, Wessel J, Gollish JD, Penney D. Assessing stability and
319 change of four performance measures: a longitudinal study evaluating outcome
320 following total hip and knee arthroplasty. *BMC Musculoskelet Disord* 2005;6:3.
321 doi:10.1186/1471-2474-6-3.
- 322 [11] Oosting E, Jans MP, Dronkers JJ, Naber RH, Dronkers-Landman CM, Appelman-de

- Vries SM, et al. Preoperative home-based physical therapy versus usual care to improve functional health of frail older adults scheduled for elective total hip arthroplasty: a pilot randomized controlled trial. *Arch Phys Med Rehabil* 2012;93:610–6. doi:10.1016/j.apmr.2011.11.006.
- [12] de Groot I, Bussmann H, Stam H, Verhaar J. Small increase of actual physical activity 6 months after total hip or knee arthroplasty. *Clin Orthop Relat Res* 2008;466:2201–8. doi:10.1007/s11999-008-0315-3.
- [13] Vissers MM, Bussmann JB, Groot IB de, Verhaar JAN, Reijman M. Physical functioning four years after total hip and knee arthroplasty. *Gait Posture* 2013;38:310–5. doi:10.1016/j.gaitpost.2012.12.007.
- [14] Harding P, Holland AE, Delany C, Hinman RS. Do activity levels increase after total hip and knee arthroplasty? *Clin Orthop Relat Res* 2014;472:1502–11. doi:10.1007/s11999-013-3427-3.
- [15] Arnold JB, Walters JL, Ferrar KE. Does Physical Activity Increase After Total Hip or Knee Arthroplasty for Osteoarthritis? A Systematic Review. *J Orthop Sport Phys Ther* 2016;46:431–42. doi:10.2519/jospt.2016.6449.
- [16] McDonald D, Siegmeth R, Deakin A, Kinninmonth AWG, Scott NB. An enhanced recovery programme for primary total knee arthroplasty in the United Kingdom-- follow up at one year. *Knee* 2012;19:525–9. doi:10.1016/j.knee.2011.07.012.
- [17] Grant PM, Ryan CG, Tigbe WW, Granat MH. The validation of a novel activity monitor in the measurement of posture and motion during everyday activities. *Br J Sports Med* 2006;40:992–7. doi:10.1136/bjsm.2006.030262.
- [18] Ryan CG, Grant PM, Tigbe WW, Granat MH. The validity and reliability of a novel activity monitor as a measure of walking. *Br J Sports Med* 2006;40:779–84. doi:10.1136/bjsm.2006.027276.
- [19] Sellers C, Dall P, Grant M, Stansfield B. Validity and reliability of the activPAL3 for measuring posture and stepping in adults and young people. *Gait Posture* 2016;43:42–7. doi:http://dx.doi.org/10.1016/j.gaitpost.2015.10.020.
- [20] Stansfield B, Hajarnis M, Sudarshan R. Characteristics of very slow stepping in healthy adults and validity of the activPAL3TM activity monitor in detecting these steps. *Med*

Eng Phys 2014. doi:10.1016/j.medengphy.2014.10.003.

[21] Söderman P, Malchau H. Is the Harris hip score system useful to study the outcome of total hip replacement? Clin Orthop Relat Res 2001;189–97.

[22] Dawson J, Fitzpatrick R, Murray D, Carr A. Questionnaire on the perceptions of patients about total knee replacement. J Bone Jt Surg 1996;78:185–90.

[23] Wylde V, Learmonth I, Cavendish V. The Oxford hip score : the patient ' s perspective. Ealth Qual Life Outcomes 2005;8:1–8. doi:10.1186/1477-7525-3-66.

[24] Rikli R, Jones J. The Reliability and Validity of a 6 Minute Walk as a measure of physical endurance in older adults. J Aging Phys Act 1998;6:363–75.

[25] O'Neill M, OO'Neill G, Perkins N. Scottish Arthroplasty Project Biennial Report 2014. 2014.

[26] Göran Garellick, Kärrholm J, Lindahl H, Malchau H, Rogmark C, Rolfson O. Swedish Hip Arthroplasty register : Annual report 2013. 2013. doi:10.1111/epp.12066.

[27] Tudor-Locke C, Craig CL, Thyfault JP, Spence JC. A step-defined sedentary lifestyle index : < 5000 steps / day. Appl Physiol , Nutr Metab 2013;114:100–14.

[28] Grant PM, Dall PM, Kerr A. Daily and hourly frequency of the sit to stand movement in older adults: a comparison of day hospital, rehabilitation ward and community living groups. Aging Clin Exp Res 2013;23:437–44. doi:10.1007/BF03325239.

[29] Taraldsen K, Askin T, Sletvold O, Einarsen E, Bjastad K, Helbostad J. Evaluation of a Body-Worn Sensor System to Measure Physical Activity in Older People With Impaired Function. vol. 91. 2011.

[30] de Groot IB, Bussmann JB, Stam HJ, Verhaar J a N. Actual everyday physical activity in patients with end-stage hip or knee osteoarthritis compared with healthy controls. Osteoarthritis Cartilage 2008;16:436–42. doi:10.1016/j.joca.2007.08.010.

[31] Vissers MM, Bussmann JB, de Groot IB, Verhaar J a N, Reijman M. Physical functioning four years after total hip and knee arthroplasty. Gait Posture 2013;38:310–5. doi:10.1016/j.gaitpost.2012.12.007.

[32] Kim WS, Kim EY. Comparing self-selected speed walking of the elderly with self-selected slow, moderate, and fast speed walking of young adults. Ann Rehabil Med

2014;38:101–8. doi:10.5535/arm.2014.38.1.101.

[33] Menz HB, Lord SR, Fitzpatrick RC. Age-related differences in walking stability. *Age Ageing* 2003;32:137–42.

[34] Tudor-Locke C, Barreira T V., Brouillette RM, Foil HC, Keller JN. Preliminary comparison of clinical and free-living measures of stepping cadence in older adults. *J Phys Act Heal* 2013;10:1175–80.

[35] Warburton DER, Nicol CW, Bredin SSD. Health benefits of physical activity: the evidence. *Can Med Assoc J* 2006;174:801–9. doi:10.1503/cmaj.051351.

[36] Fujita K, Makimoto K, Tanaka R, Mawatari M, Hotokebuchi T. Prospective study of physical activity and quality of life in Japanese women undergoing total hip arthroplasty. *J Orthop Sci* 2013;18:45–53. doi:10.1007/s00776-012-0318-5.

[37] Rolving N, Obbling KH, Christensen FB, Fonager K. Physical activity level, leisure activities and related quality of life 1 year after lumbar decompression or total hip arthroplasty. *Eur Spine J* 2013;22:802–8. doi:10.1007/s00586-012-2535-1.

Suppliers


^a **Stryker Orthopaedics**, Michigan, USA: Exeter® femoral component, Contemporary® cemented cup, Trident® uncemented cup, X3 polyethylene liner.

^b **PAL Technologies Ltd.** Glasgow, UK: activPAL3™


List of Figures

Figure 1 Strobe flow chart of participant recruitment.


Figure 2 Model outcomes for significant relationships by months-post operative. Note in figures a) to e) for illustrative purposes time and time² have been left in the models even though they were not significant (see Table 2 for significant model parameters). a) Upright time by BMI; b) Steps by BMI; c) Longest upright bout by Harris Hip Score baseline; d) Largest number of steps in an upright bout by BMI; e) Cadence of stepping bouts >60s by HHS baseline; f) Harris Hip Score by BMI; g) Harris Hip Score by age; h) Oxford Hip Score by Harris Hip Score baseline; i) Six minute walk test by BMI.




a)




f)




b)




g)




c)




h)



d)



i)



e)




Table 1 Outcomes at each time point.

Outcome	Pre-operative (Mean SD)	3m Post-operative (Mean SD)	12m Post-operative (Mean SD)
Primary outcomes			
Sit-to-stand transitions (/day)	46 (11)	44 (11)	44 (11)
Upright time (hours/day)	5.35 (2.07)	5.55 (1.74)	5.42 (1.61)
Steps (/day)	5320 (3015)	5943 (2675)	6155 (2631)
Secondary outcomes			
Longest upright bout (hours)	1.78 (1.48)	1.58 (0.82)	2.09 (1.96)
Largest number of steps in an upright bout	1934 (1480)	2559 (1841)	2671 (1705)
Cadence of bouts >60s (steps/min)	85 (16)	91 (13)	93 (12)
Harris Hip Score (/100)	50 (10)	88 (10)	91 (11)
Oxford Hip Score (/48)	15 (6)	42 (7)	44 (6)
Six minute walk test (m)	270 (93)	374 (87)	399 (104)

Table 2 Mixed linear model results, both fixed and random parameters for all outcomes. Those elements of the model that are included produced an improvement in 2LnL of a minimum of 3.84. Variables were centred as: BMI 30 kg/m², Age 70 years, HHS Baseline 50. Beta = change in outcome per unit of model element: Time (months), BMI (kg/m²), Age (years), HHS (Harris Hip Score) Baseline (score/100). 95% confidence intervals are given.

Outcome	Fixed parameters										Random parameter	
	Intercept		Time		Time ²		BMI		Age		HHS Baseline	
	Beta	P value	Beta	P value	Beta	P value	Beta	P value	Beta	P value	Beta	P value
Primary outcomes												
Sit-to-stand transitions (/day)	44.4	<0.001									75.7	0.002
	(40.8,47.9)										(40.2,142.5)	
Upright time (hours/day)	5.52	<0.001					-0.153	0.003			1.630	0.002
	(4.99,6.05)						(-0.248,-0.057)				(0.862,3.087)	
Steps (/day)	5950	<0.001					-263	0.001			3270000	0.003
	(5184,6715)						(-401,-126)				(1680000,6365000)	
Secondary outcomes												
Longest upright bout (hours)	1.81	<0.001									0.0325	0.053
	(1.50,2.12)										(-0.0004,0.0655)	
Largest number of steps in an upright bout	2453	<0.001					-144	0.001			783500	0.031
	(2023,2883)						(-221,-67)				(315500,1945300)	
Cadence of bouts >60s (steps/min)	89.1	<0.001									0.696	0.004
	(84.9,93.2)										(0.240,1.151)	
Harris Hip Score (/100)	51.3	<0.001	15.84	<0.001	-1.04	<0.001	-0.41	0.070	0.38	0.047	15.47	0.184
	(47.7,54.9)		(13.96,17.72)		(-1.18,-0.89)		(-0.85,0.04)		(0.01,0.76)		(3.53,67.70)	
Oxford Hip Score (/48)	14.9	<0.001	11.16	<0.001	-0.73	<0.001					0.19	0.029
	(12.5,17.2)		(9.99,12.33)		(-0.82,-0.64)						(-0.003,0.383)	
Six minute walk test (m)	274	<0.001	42.62	<0.001	-2.646	<0.001	-6.09	0.03			5016	0.002
	(241,307)		(30.54,54.69)		(-3.582,-1.709)		(-11.36,-0.83)				(2671,9419)	

Funding source

This study was funded by Glasgow Caledonian University.

Acknowledgements

The authors would like to thank all participants in the study and the Rehab and Orthopaedic staff of the Golden Jubilee Hospital for their support in recruitment of participant. The authors thank Dr David McDonald (Golden Jubilee National Hospital) for his contribution to the development of the research study, through the provision of advice on project development and facilitating data collection.